

Shears

$$\tilde{\mathbf{A}} = \begin{bmatrix} 1 & h_{xy} & h_{xz} & 0 \\ h_{yx} & 1 & h_{yz} & 0 \\ h_{zx} & h_{zy} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


Shears y into x

7


7

Rotations

- 3D Rotations fundamentally more complex than in 2D
 - 2D: amount of rotation
 - 3D: amount and axis of rotation



-VS-




8

8

Rotations

- Rotations still orthonormal
- $\text{Det}(\mathbf{R}) = 1 \neq -1$
- Preserve lengths and distance to origin
- 3D rotations **DO NOT COMMUTE!**
- Right-hand rule
- Unique matrices




9

9

Axis-aligned 3D Rotations

- 2D rotations implicitly rotate about a third out of plane axis



10

10

Axis-aligned 3D Rotations

- Also known as “direction-cosine” matrices

$$\mathbf{R}_x = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix} \quad \mathbf{R}_y = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$

$$\mathbf{R}_z = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

14


14

Arbitrary Rotations

- Can be built from axis-aligned matrices:

$$\mathbf{R} = \mathbf{R}_{\hat{z}} \cdot \mathbf{R}_{\hat{y}} \cdot \mathbf{R}_{\hat{x}}$$

- Result due to Euler... hence called Euler Angles
- Easy to store in vector
 $\mathbf{R} = \text{rot}(x, y, z)$
- But NOT a vector.




15

15

Exponential Maps

- Direct representation of arbitrary rotation
- AKA: axis-angle, angular displacement vector
- Rotate θ degrees about some axis
- Encode θ by length of vector

$$\theta = |\mathbf{r}|$$



18

18

Exponential Maps

- Given vector \mathbf{r} , how to get matrix \mathbf{R}
- Method from text:
 1. rotate about x axis to put \mathbf{r} into the x - y plane
 2. rotate about z axis align \mathbf{r} with the x axis
 3. rotate θ degrees about x axis
 4. undo #2 and then #1
 5. composite together


19

19

Exponential Maps

◦ Rodriguez Formula

$$\begin{aligned}\mathbf{x}' &= \hat{\mathbf{r}}(\hat{\mathbf{r}} \cdot \mathbf{x}) \\ &+ \sin(\theta)(\hat{\mathbf{r}} \times \mathbf{x}) \\ &- \cos(\theta)(\hat{\mathbf{r}} \times (\hat{\mathbf{r}} \times \mathbf{x}))\end{aligned}$$



Actually a minor variation ... 22


22

Exponential Maps

◦ Rodriguez Formula

$$\begin{aligned}\mathbf{x}' &= \hat{\mathbf{r}}(\hat{\mathbf{r}} \cdot \mathbf{x}) \\ &+ \sin(\theta)(\hat{\mathbf{r}} \times \mathbf{x}) \\ &- \cos(\theta)(\hat{\mathbf{r}} \times (\hat{\mathbf{r}} \times \mathbf{x}))\end{aligned}$$

Linear in \mathbf{x}



Actually a minor variation ... 22

22

Exponential Maps

- Building the matrix

$$\mathbf{x}' = ((\hat{\mathbf{r}}\hat{\mathbf{r}}^t) + \sin(\theta)(\hat{\mathbf{r}}\times) - \cos(\theta)(\hat{\mathbf{r}}\times)(\hat{\mathbf{r}}\times)) \mathbf{x}$$

$$(\hat{\mathbf{r}}\times) = \begin{bmatrix} 0 & -\hat{r}_z & \hat{r}_y \\ \hat{r}_z & 0 & -\hat{r}_x \\ -\hat{r}_y & \hat{r}_x & 0 \end{bmatrix}$$

Antisymmetric matrix

$(\mathbf{a}\times)\mathbf{b} = \mathbf{a}\times\mathbf{b}$

Easy to verify by expansion

23

23

Exponential Maps

- Allows tumbling
- No gimbal-lock!
- Orientations are space within π -radius ball
- Nearly unique representation
- Singularities on shells at 2π
- Nice for interpolation

24

24

Quaternions

- Multiplication natural consequence of defn.

$$q \cdot p = (\mathbf{z}_q s_p + \mathbf{z}_p s_q + \mathbf{z}_p \times \mathbf{z}_q, s_p s_q - \mathbf{z}_p \cdot \mathbf{z}_q)$$

- Conjugate

$$q^* = (-\mathbf{z}, s)$$

- Magnitude

$$\|q\|^2 = \mathbf{z} \cdot \mathbf{z} + s^2 = q \cdot q^*$$

31

31

Quaternions

- Vectors as quaternions

$$v = (\mathbf{v}, 0)$$

- Rotations as quaternions

$$r = \left(\hat{\mathbf{r}} \sin \frac{\theta}{2}, \cos \frac{\theta}{2} \right)$$

- Rotating a vector

$$x' = r \cdot x \cdot r^* \quad \leftarrow \text{Compare to Exp. Map}$$

- Composing rotations

$$r = r_1 \cdot r_2$$

32

32


Quaternions

- No tumbling
- No gimbal-lock
- Orientations are “double unique”
- Surface of a 3-sphere in 4D $||r|| = 1$
- Nice for interpolation

33

33

Interpolation



34

34
